



Online Version ISSN: 1314-412X
Volume 3, Number 4
December 2011

AGRICULTURAL SCIENCE AND TECHNOLOGY

2011

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Editor-in-Chief

Tsanko Yablanski
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Co-Editor-in-Chief

Radoslav Slavov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Editors and Sections

Genetics and Breeding

Atanas Atanassov (Bulgaria)
Ihsan Soysal (Turkey)
Max Rothschild (USA)
Stoitcho Metodiev (Bulgaria)

Nutrition and Physiology

Nikolai Todorov (Bulgaria)
Peter Surai (UK)
Zervas Georgios (Greece)

Production Systems

Dimitar Pavlov (Bulgaria)
Dimitar Panaiotov (Bulgaria)
Jordan Staikov (Bulgaria)
Georgi Zhelyazkov (Bulgaria)

Agriculture and Environment

Georgi Petkov (Bulgaria)
Ramesh Kanwar (USA)

Product Quality and Safety

Marin Kabakchiev (Bulgaria)
Stefan Denev (Bulgaria)

English Editor

Yanka Ivanova (Bulgaria)

Scope and policy of the journal

Agricultural Science and Technology /AST/ – an International Scientific Journal of Agricultural and Technology Sciences is published in English in one volume of 4 issues per year, as a printed journal and in electronic form. The policy of the journal is to publish original papers, reviews and short communications covering the aspects of agriculture related with life sciences and modern technologies. It will offer opportunities to address the global needs relating to food and environment, health, exploit the technology to provide innovative products and sustainable development. Papers will be considered in aspects of both fundamental and applied science in the areas of Genetics and Breeding, Nutrition and Physiology, Production Systems, Agriculture and Environment and Product Quality and Safety. Other categories closely related to the above topics could be considered by the editors. The detailed information of the journal is available at the website. Proceedings of scientific meetings and conference reports will be considered for special issues.

Submission of Manuscripts

All manuscript written in English should be submitted as MS-Word file attachments via e-mail to ascitech@uni-sz.bg. Manuscripts must be prepared strictly in accordance with the detailed instructions for authors at the website <http://www.uni-sz.bg/ascitech/index.html> and the instructions on the last page of the journal. For each manuscript the signatures of all authors are needed confirming their consent to publish it and to nominate an author for correspondence. They have to be presented by a submission letter signed by all authors. The form of the submission letter is available upon request from the Technical Assistance or could be downloaded from the website of the journal. All manuscripts are subject to editorial review and the editors reserve the right to improve style and return the paper for rewriting to the authors, if necessary. The editorial board reserves rights to reject manuscripts based on priorities and space availability in the journal.

Subscriptions

Agricultural Science and Technology is published four times a year. The subscription price for institutions is 80 € and for personal subscription 30 € which

include electronic access and delivery. Subscription run for full calendar year. Orders, which must be accompanied by payment may be sent direct to the publisher:

Trakia University
Faculty of Agriculture, Bank account:
UniCredit Bulbank,
Sofia BIC: UNCRBGSF

IBAN: BG29UNCR76303100117681
With UniCredit Bulbank Stara Zagora

Internet Access

This journal is included in the Trakia University Journals online Service which can be found at www.uni-sz.bg.

Copyright


All rights reserved. No part of this publications may be translated into other languages, reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying or any information storage and retrieval system without permission in writing from the publisher.

Address of Editorial office:

Agricultural Science and Technology
Faculty of Agriculture, Trakia University
Student's campus, 6000 Stara Zagora
Bulgaria
Telephone.: +359 42 699330
+359 42 699446
<http://www.uni-sz.bg/ascitech/index.html>

Technical Assistance:

Nely Tzvetanova
Telephone.: +359 42 699446
E-mail: ascitech@uni-sz.bg



*AGRICULTURAL
SCIENCE AND TECHNOLOGY*

2011

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Effect of the rhizobacterium *Bacillus subtilis* on the development of the root-knot nematode *Meloidogyne arenaria* at different temperatures

M. Mohamedova*, H. Samaliev

Department of Entomology, Faculty of Plant Protection and Agroecology, Agricultural University, 12 Mendeleev, 4000 Plovdiv, Bulgaria

Abstract. Growth room experiment was carried out to investigate the effect of different temperatures on the efficacy of the rhizobacterium *Bacillus subtilis* on the development of the root-knot nematode *Meloidogyne arenaria*. Potato plants (*Solanum tuberosum* L.) were inoculated with *M. arenaria* second-stage juveniles. At 7th, 14th and 21th days after nematode inoculation the rhizosphere of the plants were treated with bacterial suspensions of the two local strains of *B. subtilis* (A1 and B1). The potato plants were grown in growth room at 16±1, 20±1, 24±1, 28±1 and 32±1°C. The rate of *M. arenaria* development in potato roots treated with *B. subtilis* (strains A1 and B1) was lower than that in the untreated plant roots. The highest efficacy of the bacterium on the development of *M. arenaria* was observed at 24-32°C.

Keywords: root-knot nematode, *Bacillus subtilis*, temperature influence, development, biological control

Introduction

Nematodes cause about 20.6% worldwide yield loss (Sasser, 1989). Root-knot nematodes (*Meloidogyne* sp.) are serious plant pathogens which parasitize on vegetables growing on fields and in greenhouses in Bulgaria (Samaliev and Stoyanov, 2008). The control of these pathogens includes mainly the use of pesticides. However, in many cases, high-toxic nematicides are required to reduce damage and increase yield (Sikora and Fernandez, 2005). Recently interest in biological control of nematodes has increased, fuelled by public concern over the use of chemicals in the environment (Moens et al., 2004; Gowen et al., 2005).

Bacillus is one of the important bacterial genera, which can suppress nematode invasion (Kloepper and Ryu, 2006). Gokte and Swarup (1988) reported that *B. subtilis*, *B. cereus* and *B. pumilus* exhibited larvicidal activity against the second stage juveniles (J2) of *Meloidogyne incognita* *in vitro*. In pot experiment Gautam et al. (1995) observed reduction of *M. incognita* multiplication on tomato due to use of *B. subtilis* as seeds treatment. *B. thuringiensis* was reported as a potential control agents to many plant parasitic nematodes, including *Meloidogyne* sp. (Borgonie et al., 1996; Marroquin et al., 2000; Khyami-Horani and Al-Banna, 2006). In our previous studies some local strains of the rhizobacterium *B. subtilis* caused mortality of *M. arenaria* J2 in laboratory experiments (Mohamedova and Samaliev, 2006) and prevented invasion of J2 into tomato roots in pot experiments at 25°C (Mohamedova and Samaliev, 2010). One of the factors limiting the effect of the rhizobacteria on root-knot nematodes is the temperature. Becker et al. (1992) reported the activity of *B. thuringiensis* var. *israelensis* towards different plant pathogens did not change significantly between 19 and 33°C. However, the effect of *Pseudomonas oryzae* on *Globodera rostochiensis* hatch and J2 is highest between 21 and 25°C and decreasing below 20°C (Andreoglou and Gowen, 2000). The objective of this evaluation was to determine the effect of the rhizobacterium *B. subtilis* on *M. arenaria* development at different temperatures.

Materials and methods

Nematode culture and sterilization

The root-knot nematode *M. arenaria* was obtained from cultures derived from single egg masses maintained on tomato (*Lycopersicon esculentum* Mill., cv. Velositi) in greenhouse at 24-26°C. Mature egg masses of *M. arenaria* were hand picked, using sterilized needle and forceps from heavily infested tomato roots. These egg masses were sterilized in streptomycin sulphate (0.1%) for 45 min (Sawhney and Webster, 1975) and rinsed in sterile distilled water (SDW) before use in experiments. J2 of *M. arenaria* were extracted from infested tomato roots. Galled roots with egg masses were washed free of soil, cut into small pieces, placed in 1.5% NaOCl and macerated in a blender (Hussey and Barker, 1973). The suspension was poured onto cotton-wool filter, incubated at 24-26°C and hatched J2 were collected every 24 h. J2 were sterilized in streptomycin sulphate (0.1%) for 15 min (Mountain, 1955) and rinsed in SDW before use in experiments.

Bacterium culture and identification

The strains B1 and A1 of *B. subtilis* were cultivated at 28°C on tryptic soy broth (TSB). For inoculum production single colonies of the two strains were grown at 27°C for 48 h with shaking (150 rpm) in 250 ml Erlenmeyer flasks containing 100 ml of TSB. The bacterial suspensions were centrifuged at 2800 g for 20 min. The concentrated suspensions were diluted with sterile tap water to give the concentrations required for experiments. Bacterial concentrations were determined by using a spectrophotometer. Bacterial identity of both strains was determined by Fatty Acid Analysis (Microbial Identification System Inc., Delaware, USA).

Effect of Bacillus subtilis (strains A1 and B1) on the development of Meloidogyne arenaria at different temperatures

Sterile potato tubers (cv. Nadezda) with one sprout were planted in closed plastic containers (4.5 cm high, 10 cm in diameter) in 200 g of sterilized soil (3:1 loam/sand mixture), at 40% moisture-

* e-mail: m.mohamedova@au-plovdiv.bg

level. This technique was modified by Phillips et al. (1980). The containers were stored in the dark at a constant temperature of 21°C for 2 weeks. Two-week-old potato plants were inoculated with 200 J2 of *M. arenaria* and placed in an incubator, in dark, at 16°C, 20°C, 24°C, 28°C and 32°C. After 7 days the plants were uprooted, washed free of soil and replanted in sterilized soil in new containers. At 7, 14 and 21 days after J2 inoculation, 20 ml of a bacterial suspension containing 10^8 cells/ml of *B. subtilis* (strains A1 and B1) were applied to the soil around the roots. Four untreated plants received 20 ml of sterile distilled water and served as control. At each treatment 4 plants (fixed) were harvested to determine the stage of nematode development when the bacterium was applied. After each treatment the plants were placed back in the incubators. At 83, 51, 42, 33 and 36 days after J2 inoculation at 16°C, 20°C, 24°C, 28°C and 32°C, respectively, the roots were collected, washed free of soil, stained in acid fuchsin (Bridge et al., 1982), macerated in a blender and the number of different stages of the nematode determined. The experiment was conducted with four replication per treatment.

Statistical analysis

All statistical analysis was performed using procedures of the SPSS programme, significance being determined at $P=0.05$. Because most of the data were counts out of a total, e.g. nematodes in roots after plant inoculation with 200 J2, most of the analysis that was done was logistic regression based on binomial data. However, some data were not analysed, because they were extreme (e.g.

almost 0 and 100%) and the differences were obvious.

Results and discussion

The effect of *B. subtilis* (strains A1 and B1) on the development and fecundity of *M. arenaria* in potato roots was different at different temperatures. The influence of the strains was more pronounced the closer the bacterium was applied after J2 inoculation and the higher the temperature was. At $28\pm 1^\circ\text{C}$ and 7 days after J2 inoculation strain B1 caused the highest inhibition of nematode development into plant roots. At this temperature only 25% of developed females had egg masses (Table 4). The effect of the strain was less marked at $24\pm 1^\circ\text{C}$ and $32\pm 1^\circ\text{C}$. However, the percentage of females with egg masses in treated plants was significantly lower than this one in untreated roots, respectively, 37.5% and 41.7% (Tables 3 and 5).

After 14 days exposure and temperatures of $24\pm 1^\circ\text{C}$, $28\pm 1^\circ\text{C}$ and $32\pm 1^\circ\text{C}$ strain B1 significantly delayed egg masses formation compared to the untreated plants. The percentage of developed females with egg masses was respectively, 46.7%, 50.0% and 63.2% (Tables 3, 4 and 5). The number of eggs per egg masses was significantly lower than in the untreated plants, when the potato plants treated with the strain B1 up to 14 days after J2 inoculation at all investigated temperatures, and even after 21 days at $24\pm 1^\circ\text{C}$, $28\pm 1^\circ\text{C}$ and $32\pm 1^\circ\text{C}$ (Tables 3, 4 and 5).

Table 1. Effect of *B. subtilis* (A1 and B1) on the different stages of development of *M. arenaria* in potato roots 83 days after inoculation with 200 J2 at $16\pm 1^\circ\text{C}$. Percentages were calculated as % of the nematodes counted in the roots

Strains	Days of bacteria treatment after J2 inoculation	Treatments	Developmental stages of <i>M. arenaria</i>				Eggs/egg mass ¹
			J2 (%) ²	J3 (%) ²	J4 (%) ²	Females with egg masses (%) ²	
A1	7	Fixed	12.5(± 4.7)	13.3(± 4.6)	74.2(± 4.1)	91.7(± 5.6)	132
			92.3	7.6	0	-	-
	14	Fixed	6.5(± 3.9)	14.4(± 4.0)	79.1(± 5.2)	100(± 0.0)	154
			37.7	62.3	0	-	-
	21	Fixed	0(± 0.0)	0(± 0.0)	100(± 0.0)	100(± 0.0)	165
			10.2	31.1	58.7	-	-
Control		0(± 0.0)	0(± 0.0)	100(± 0.0)	100(± 0.0)	168	
s.e.d						14.3	
B1	7	Fixed	24.6(± 4.8)	23.3(± 4.7)	52.1(± 5.4)	85.0(± 6.1)	138
			96.7	3.3	0	-	-
	14	Fixed	32.0(± 6.2)	18.2(± 5.3)	49.8(± 5.1)	95.5(± 6.4)	155
			44.5	55.5	0	-	-
	21	Fixed	0(± 0.0)	0(± 0.0)	100(± 0.0)	100(± 0.0)	174
			6.4	32.7	60.9	-	-
Control		0(± 0.0)	0(± 0.0)	100(± 0.0)	100(± 0.0)	178	
s.e.d						17.8	

¹ Data were analysed by analysis of variance. Summary statistics are mean values and the standard error of the difference.

² Data were analysed using logistic regression analysis. Summary statistics are mean counts (\pm standard error).

³ Fixed plants were not analysed.

Table 2. Effect of *B. subtilis* (A1 and B1) on the different stages of development of *M. arenaria* in potato roots 51 days after inoculation with 200 J2 at 20±1°C. Percentages were calculated as % of the nematodes counted in the roots

Strains	Days of bacteria treatment after J2 inoculation	Treatments	Developmental stages of <i>M. arenaria</i>				Eggs/egg mass ¹
			J2 (%) ²	J3 (%) ²	J4 (%) ²	Females with egg masses (%) ²	
A1	7	Fixed	10.7(±6.8)	11.3(±4.4)	78.0(±5.7)	94.7(±3.9)	128
			89.8	10.2	0	-	-
	14	Fixed	3.9(±4.0)	11.5(±4.8)	84.6(±6.1)	95.5(±4.7)	160
			28.6	71.4	0	-	-
	21	Fixed	0(±0.0)	0(±0.0)	100(±0.0)	100(±0.0)	164
			1.3	19.5	79.2	-	-
Control		0(±0.0)	0(±0.0)	100(±0.0)	100(±0.0)	176	
s.e.d						15.8	
B1	7	Fixed	15.7(±4.8)	16.5(±5.2)	67.8(±6.1)	72.7(±6.7)	119
			92.2	7.8	0	-	-
	14	Fixed	12.8(±6.0)	14.8(±6.6)	72.4(±6.5)	83.3(±5.8)	147
			24.5	75.5	0	-	-
	21	Fixed	0(±0.0)	0(±0.0)	100(±0.0)	100(±0.0)	160
			1.1	17.2	81.7	-	-
Control		0(±0.0)	0(±0.0)	100(±0.0)	100(±0.0)	171	
s.e.d						19.4	

Table 3. Effect of *B. subtilis* (A1 and B1) on the different stages of development of *M. arenaria* in potato roots 42 days after inoculation with 200 J2 at 24±1°C. Percentages were calculated as % of the nematodes counted in the roots

Strains	Days of bacteria treatment after J2 inoculation	Treatments	Developmental stages of <i>M. arenaria</i>				Eggs/egg mass ¹
			J2 (%) ²	J3 (%) ²	J4 (%) ²	Females with egg masses (%) ²	
A1	7	Fixed	5.8(±5.9)	11.4(±3.6)	82.8(±5.7)	37.5(±6.5)	81
			87.3	12.7	0	-	-
	14	Fixed	0(±0.0)	6.8(±4.9)	93.2(±3.8)	46.7(±6.1)	93
			10.7	86.2	3.1	-	-
	21	Fixed	0(±0.0)	0(±0.0)	100(±0.0)	50.0(±5.2)	125
			0	8.5	91.5	-	-
Control		0(±0.0)	0(±0.0)	100(±0.0)	100(±0.0)	183	
s.e.d						17.1	
B1	7	Fixed	6.8(±3.9)	7.6(±4.1)	85.6(±3.8)	68.2(±5.5)	122
			81.2	18.8	0	-	-
	14	Fixed	1.1(±3.4)	3.5(±6.7)	95.4(±3.3)	81.5(±4.9)	165
			12.2	82.0	5.8	-	-
	21	Fixed	0(±0.0)	0(±0.0)	100(±0.0)	100(±0.0)	174
			2.8	7.8	89.4	-	-
Control		0(±0.0)	0(±0.0)	100(±0.0)	100(±0.0)	189	
s.e.d						18.5	

Table 4. Effect of *B. subtilis* (A1 and B1) on the different stages of development of *M. arenaria* in potato roots 33 days after inoculation with 200 J2 at 28±1°C. Percentages were calculated as % of the nematodes counted in the roots

Strains	Days of bacteria treatment after J2 inoculation	Treatments	Developmental stages of <i>M. arenaria</i>				Eggs/egg mass ¹
			J2 (%) ²	J3 (%) ²	J4 (%) ²	Females with egg masses (%) ²	
A1	7	Fixed	4.6(±5.3)	15.4(±6.7)	80.0(±4.1)	25.0(±6.4)	84
			84.2	15.8	0	-	-
	14	Fixed	0(±0.0)	4.8(±6.9)	95.2(±5.0)	50.0(±5.7)	99
			8.9	87.4	3.7	-	-
	21	Fixed	0(±0.0)	0(±0.0)	100(±0.0)	47.1(±6.0)	126
			0	5.1	94.9	-	-
Control		0(±0.0)	0(±0.0)	100(±0.0)	100(±0.0)	186	
s.e.d						25.5	
B1	7	Fixed	3.3(±5.8)	11.7(±4.7)	85.0(±5.1)	72.2(±4.6)	124
			80.2	19.8	0	-	-
	14	Fixed	0(±0.0)	3.0(±3.7)	97.0(±5.0)	88.0(±3.9)	166
			9.6	85.2	5.2	-	-
	21	Fixed	0(±0.0)	0(±0.0)	100(±0.0)	100(±0.0)	175
			0	7.3	92.7	-	-
Control		0(±0.0)	0(±0.0)	100(±0.0)	100(±0.0)	190	
s.e.d						17.2	

Table 5. Effect of *B. subtilis* (A1 and B1) on the different stages of development of *M. arenaria* in potato roots 36 days after inoculation with 200 J2 at 32±1°C. Percentages were calculated as % of the nematodes counted in the roots

Strains	Days of bacteria treatment after J2 inoculation	Treatments	Developmental stages of <i>M. arenaria</i>				Eggs/egg mass ¹
			J2 (%) ²	J3 (%) ²	J4 (%) ²	Females with egg masses (%) ²	
A1	7	Fixed	1.9(±3.5)	7.2(±5.8)	90.9(±5.6)	41.7(±6.8)	92
			79.2	20.8	0	-	-
	14	Fixed	0(±0.0)	6.1(±6.0)	93.9(±4.7)	63.2(±6.3)	103
			3.4	89.8	6.8	-	-
	21	Fixed	0(±0.0)	0(±0.0)	100(±0.0)	80.6(±6.0)	142
			0	3.2	96.8	-	-
Control		0(±0.0)	0(±0.0)	100(±0.0)	100(±0.0)	180	
s.e.d						22.4	
B1	7	Fixed	1.3(±4.2)	4.8(±3.7)	93.9(±3.8)	88.0(±4.3)	120
			77.7	22.3	0	-	-
	14	Fixed	0(±0.0)	1.6(±3.0)	98.4(±5.1)	93.3(±4.4)	155
			4.4	87.1	8.5	-	-
	21	Fixed	0(±0.0)	0(±0.0)	100(±0.0)	100(±0.0)	168
			0	4.1	95.9	-	-
Control		0(±0.0)	0(±0.0)	100(±0.0)	100(±0.0)	188	
s.e.d						19.7	

B. subtilis (strain A1) showed lower effect on the development and reproduction of *M. arenaria* into plant roots than the strain B1 (Tables 1, 2, 3, 4 and 5). Significantly lower percentage of females with egg masses into roots treated with the strain A1 compared to untreated plants was observed only after one day exposure and temperatures of 24±1°C and 28±1°C, respectively, 68.2% and 72.2% (Tables 3 and 4).

This study demonstrates that the bacterium *B. subtilis* (strain B1) inhibits *M. arenaria* development and fecundity into potato roots. In the bioassays the effect of the strain depended on the temperature. The best effectiveness of the B1 bacterial suspension was observed at temperatures from 24°C to 28°C and in the cases when the plant roots contained J2 and J3 at the time of strain application. The nematicidal effect appears quickly at 24±1°C, 28±1°C and 32±1°C and slowly at 20±1°C. There was no effect of B1 at 16±1°C. Our results are consistent with this reported by Andreoglou and Gowen (2000). The authors observed the highest effect of *P. oryzihabitans* against *G. rostochiensis* at temperatures from 22°C to 26°C. Hackenberg and Sikora (1994) reported that the rhizobacterium *Agrobacterium radiobacter* reduced the development of *G. pallida* significantly up to 70%, both at 20°C and 25°C.

B. subtilis (strain A) caused 90% mortality of *M. arenaria* J2 and significantly decreased egg hatching *in vitro* at temperatures between 24°C and 30°C. However its effect *in vivo* was insignificant. This probably due to delayed and less bacterial colonization around plant roots or restricted possibility to produce secondary metabolites which are released into the soil and interfere with *M. arenaria* at they inhibit its development. Similarly Siddiqui and Mahmood (1992) reported that two strains of *B. licheniformis* with the same effect against *M. incognita* *in vitro* showed different effectiveness *in vivo*. The authors explained the results with different levels of root colonization of the strains.

Conclusion

Our study indicates that the rhizobacterium *B. subtilis* (strain B1) is nematicidal *in vivo* at temperatures between 20°C and 32°C. Below and above these temperatures the effect of the strain on the development and reproduction of *M. arenaria* in plant roots is insignificant. Usually planting of the vegetables (tomato) in plastic greenhouses occurs at the beginning of March, when the invasive J2 of *M. arenaria* infect plant roots at soil temperature of below 16°C, i. e. lower that effective ones for *B. subtilis* (strain B1) used in this experiment. However, egg laying in the egg masses starts at soil temperatures of over 17°C In this early periods of plant growth is recommend to combine the application of the strain B1 with reduced dosage of nonfumigant nematocides to achieve successful control of *M. arenaria*.

After all the selection of flexible strains effective at lower temperatures could be an important alternative in the development of *B. subtilis* as a biological control agent.

References

Andreoglou FI and Gowen SR, 2000. Temperature response of the bacterium *Pseudomonas oryzihabitans*, from the entomopathogenic nematode *Steinernema abbasi*, on the potato cyst nematode *Globodera rostochiensis*. *Aspects of Applied Biology*, 59, 67-73.

Borgonie G, Claeys M, Leyns G, Arnaut G, Waele De D and Coomans A, 1996. Effect of nematicidal *Bacillus thuringiensis* strains on free living nematodes. 1. Light microscopic observation, spwcies and biological stage specificity and identification of resistant mutants of *Caenorhabditis elegans*. *Fundamental and Applied Nematology*, 19, 391-398.

Becker N, Zgonib, M, Ludwig M, Petric D. and Rettich F, 1992. Factors influencing the activity of *Bacillus thuringiensis* var. *israelensis* treatments. *Journal of the American Mosquito Control Association*, 8, 285-289.

Bridge J, Page S and Jordan S, 1982. An improved method for staining nematodes in roots. In: *Nematology Department, Rothamsted Report for 1981*, Harpenden, UK, 1, 171.

Gautam A, Siddiqui I and Mahmood A, 1995. Integrated management of *Meloidogyne incognita* on tomato. *Nematology of the Mediterterranean*, 23, 245-272.

Gokte N and Swarup G, 1988. On the potential of some bacterial biocides against root-knot and cyst nematodes. *Indian Journal of Nematology*, 18, 152-153.

Gowen S, Queneherve P, Fogian R, 2005. Nematode parasites of bananas and plantains. In: *Plant parasitic nematodes in subtropical and tropical agriculture*, eds. Luc, M., R. Sikora, J. Bridge, Wallingford, UK, CAB International, 611-643.

Hackenberg C and Sikora R.A, 1994. Influence of temperature and soil moisture on the biological control of the potato-cyst-nematode *Globodera pallida* using the plant-health-promoting rhizobacterium *Agrobacterium radiobacter*. *Journal of Phytopathology*, 142, 338-344.

Hussey RS, Berker KR, 1973. A comparison of methods of collecting inocula of *Meloidogyne* spp. including a new technique. *Plant Disease Reporter*, 57, 1025-1028.

Khyami-Horani H and Al-Banna L, 2006. Efficacy of *Bacillus thuringiensis jordanica* against *Meloidogyne javanica* infecting tomato. *Phytopathologia Mediterranea*, 45, 153-157.

Kloepper J and Ryu C, 2006. Bacterial endophytes as elicitors of induced systemic resistance. In: *Microbial root endophytes*, eds. Schulz, B., C. Boyle, T. Sieber. Springer-Verlag, Heildelberg, 33-51.

Marroquin LD, Elyassnia JS, Griffiths JS, Feitelson JS and Aroian RV, 2000. *Bacillus thuringiensis* (Bt) toxin susceptibility and isolation of resistance mutants in the nematode *Caenorhabditis elegans*. *Genetics*, 155, 1693-1699

Moens T, Araya M, Swennen R and Waele D De, 2004. Enhanced biodegradation of nematicides after repetitive application and its effect on root and yield parameters in commercial banana plantations. *Biology and fertility of soils*, 39, 407-414 .

Mohamedova M and Samaliev H, 2006. The effects of the rhizobacteria *Bacillus* sp. and *Pseudomonas* sp. on the root-knot nematode *Meloidogyne arenaria*. *Experimental Pathology and Parasitology*, 9, 3, 10-15.

Mohamedova M and Samaliev H, 2010. Influence of the rhizobacterium *Bacillus subtilis* on the root-knot nematode *Meloidogyne arenaria*, Jubilee scientific conference on Tradition and Challenges of Agricultural Education, Science and Business, 14-17 October. Academic Publishing House of the Agricultural University, LV, 2, 155-164.

Mountain W, 1955. A method of culturing plant-parasitic nematodes under sterile conditions. *Proceedings of the Helminthological Society of Washington*, 16, 154-155.

Philips MS, Forest JMS and Wilson LA, 1980. Screening for resistance to potato cyst nematode using closed containers. *Annals*

of Applied Biology, 6, 317-322.

Samaliev H and Stoyanov D, 2008. Parasitic nematodes of crop plants and their control. Agriculture University Academic Press, Plovdiv, 78-122.

Sasser J, 1989. Plant parasitic nematodes: the farmer's hidden enemy. A cooperative publication of the Department of Plant Pathology and Consortium for International Crop Protection, 115.

Sawhney, R and Webster JM, 1975. The role of plant growth hormones in determining the resistance of tomato plants to the root-

knot nematode, *Meloidogyne incognita*. Nematologica, 21, 95-103.

Siddiqui ZA and Mahmood I, 1992. Biological control of root-knot disease complex of chickpea caused by *Meloidogyne incognita* race 3 and *Macrophomina phaseolina*. Nematologia Mediterranea, 20, 199-202.

Sikora R and Fernandez E, 2005. Nematode parasites of vegetables. In: Plant parasitic nematodes in subtropical and tropical agriculture, eds. Luc, M., R. Sikora, J. Bridge, Wallingford, UK. CAB International, 319-392.

CONTENTS

1 / 2

Reviews

- 1. Lameness scoring systems for cattle in dairy farms** 291
T. Penev

Genetics and Breeding

- 2. Body and carcass parameters of sea bream (*Sparus aurata* L.) and sea bass (*Dicentrarchus labrax* L.)** 299
M. Marinova, I. Sirakov, Y. Staykov, E. Ivancheva
- 3. Effect of breed upon blood lysozyme and complement activity in different sheep breeds** 302
L. Sotirov, Ts. Koynarski, V. Semerdjiev, D. Dimov, S. Laleva, P. Slavova, M. Iliev, D. Yarkov
- 4. Winter wheat productivity under favorable and drought environments III. Effect of fertilization** 306
A. Ivanova, N. Tsenov
- 5. Evaluation of perspective sorghum breeding forms in their reaction to some diseases in field conditions** 310
M. Georgieva – Andreeva, K. Tanova, S. Raykov

Nutrition and Physiology

- Effect of dietary coconut oil supplementation on some blood biochemical indices in yearling rams** 313
T. Slavov, V. Radev, K. Sivkova, I. Varlyakov
- Pharmacokinetics of tilmicosin after oral application of Pulmotil G 200 – premix in pigs** 318
D. Dimitrova, V. Katsarov, D. Dimitrov, D. Tsoneva
- Epidermal growth factor content in rabbit doe milk during the different lactation stages** 323
E. Vachkova, B. Bivolarski
- Intraorbital glands in turkey broilers. III. Lacrimal gland histometry** 327
D. Dimitrov
- Effect of body condition score at calving on body condition during lactation in Holstein and Brown Swiss cows** 330
Zh. Gergovska, T. Angelova, D. Yordanova, Zh. Krastanov, Ch. Miteva
- Use of brewer's grains for feeding of lambs** 336
A. Kirilov, K. Ivanov

Production Systems

- The effect of the milking liner design on the parameters of the milking machine pulsation system** 339
V. Vlashev, B. Banev, K. Peichev, G. Dineva
- Accumulation dynamic of *Ruta graveolens* L. essential oil** 343
A. Dzurmanski, G. Zhekova, D. Angelova
- Research on the water regimen of soil upon the production of vine planting material** 346
N. Kovachev, N. Taneva, V. Kovachev, L. Halil

Agriculture and Environment

- Influence of the farming, soil cultivation and Fertilization on the yield of wheat** 351
M. Nankov, L. Glogova
- Study on the applicability of a natural geomaterial for mononitrophenol removal from simulated agricultural run-off water** 354
Z. Yaneva, B. Koumanova, N. Georgieva

CONTENTS

2 / 2

Comparative technical and economic analysis of variants for cleaning and storage of manure on a farm for 108 – 120 dairy cows	359
V. Dimova, D. Dinev, Y. Popova, Y. Mitev	
Distribution of the black mussel <i>Mytilus galloprovincialis</i> (L.) along the Bulgarian Black Sea coast	368
E. Petrova, St. Stoykov	
Toxicity of plant protection products towards the imago of <i>Encarsia Formosa</i> Gah.	374
V. Yankova, S. Masheva, B. Boev, K. Toskov	
Effect of the rhizobacterium <i>Bacillus subtilis</i> on the development of the root-knot nematode <i>Meloidogyne arenaria</i> at different temperatures	378
M. Mohamedova, H. Samaliev	
Product Quality and Safety	
<hr/>	
Fatty acid composition of yogurt supplemented with walnut extract	384
S. Boycheva, N. Naydenova, G. Mihaylova, T. Dimitrov, D. Pavlov	
Near Infrared Spectroscopy for monitoring changes during yellow cheese ripening	390
S. Atanassova, N. Naydenova, T. Kolev, T. Iliev, G. Mihaylova	
Short communications	
<hr/>	
Mechanical correction the traction weight of a farm wheeled tractor	395
D. Irinchev	
Possibilities for increasing the yield and quality of asters (<i>Callistephus chinensis</i> L.) cut flower	397
N. Miteva, O. Tafradziiski	

Instruction for authors

Preparation of papers

Papers shall be submitted at the editorial office typed on standard typing pages (A4, 30 lines per page, 62 characters per line). The editors recommend up to 15 pages for full research paper (including abstract, references, tables, figures and other appendices)

The manuscript should be structured as follows: Title, Names of authors and affiliation address, Abstract, List of keywords, Introduction, Material and methods, Results, Discussion, Conclusion, Acknowledgements (if any), References, Tables, Figures.

The title needs to be as concise and informative about the nature of research. It should be written with small letter /bold, 14/ without any abbreviations.

Names and affiliation of authors

The names of the authors should be presented from the initials of first names followed by the family names. The complete address and name of the institution should be stated next. The affiliation of authors are designated by different signs. For the author who is going to be corresponding by the editorial board and readers, an E-mail address and telephone number should be presented as footnote on the first page. Corresponding author is indicated with *.

Abstract should be not more than 350 words. It should be clearly stated what new findings have been made in the course of research. Abbreviations and references to authors are inadmissible in the summary. It should be understandable without having read the paper and should be in one paragraph.

Keywords: Up to maximum of 5 keywords should be selected not repeating the title but giving the essence of study.

The introduction must answer the following questions: What is known and what is new on the studied issue? What necessitated the research problem, described in the paper? What is your hypothesis and goal?

Material and methods: The objects of research, organization of experiments, chemical analyses, statistical and other methods and conditions applied for the experiments should be described in detail. A criterion of sufficient information is to be

possible for others to repeat the experiment in order to verify results.

Results are presented in understandable tables and figures, accompanied by the statistical parameters needed for the evaluation. Data from tables and figures should not be repeated in the text.

Tables should be as simple and as few as possible. Each table should have its own explanatory title and to be typed on a separate page. They should be outside the main body of the text and an indication should be given where it should be inserted.

Figures should be sharp with good contrast and rendition. Graphic materials should be preferred. Photographs to be appropriate for printing. Illustrations are supplied in colour as an exception after special agreement with the editorial board and possible payment of extra costs. The figures are to be each in a single file and their location should be given within the text.

Discussion: The objective of this section is to indicate the scientific significance of the study. By comparing the results and conclusions of other scientists the contribution of the study for expanding or modifying existing knowledge is pointed out clearly and convincingly to the reader.

Conclusion: The most important consequences for the science and practice resulting from the conducted research should be summarized in a few sentences. The conclusions shouldn't be numbered and no new paragraphs be used. Contributions are the core of conclusions.

References:

In the text, references should be cited as follows: single author: Sandberg (2002); two authors: Andersson and Georges (2004); more than two authors: Andersson et al.(2003). When several references are cited simultaneously, they should be ranked by chronological order e.g.: (Sandberg, 2002; Andersson et al., 2003; Andersson and Georges, 2004). References are arranged alphabetically by the name of the first author. If an author is cited more than once, first his individual publications are given ranked by year, then come publications with one co-author, two co-authors, etc. The names of authors, article and journal titles in the Cyrillic or alphabet different from Latin, should be transliterated into Latin and article titles should be translated into English. The original language of articles and books translated into English is indicated in

parenthesis after the bibliographic reference (Bulgarian = Bg, Russian = Ru, Serbian = Sr, if in the Cyrillic, Mongolian = Mo, Greek = Gr, Georgian = Geor., Japanese = Ja, Chinese = Ch, Arabic = Ar, etc.)

The following order in the reference list is recommended:

Journal articles: Author(s) surname and initials, year. Title. Full title of the journal, volume, pages. Example:

Simm G, Lewis RM, Grundy B and Dingwall WS, 2002. Responses to selection for lean growth in sheep. *Animal Science*, 74, 39-50

Books: Author(s) surname and initials, year. Title. Edition, name of publisher, place of publication. Example: **Oldenbroek JK**, 1999. Genebanks and the conservation of farm animal genetic resources, Second edition. DLO Institute for Animal Science and Health, Netherlands.

Book chapter or conference proceedings: Author(s) surname and initials, year. Title. In: Title of the book or of the proceedings followed by the editor(s), volume, pages. Name of publisher, place of publication. Example:

Mauff G, Pulverer G, Operkuch W, Hummel K and Hidden C, 1995. C3-variants and diverse phenotypes of unconverted and converted C3. In: *Provides of the Biological Fluids* (ed. H. Peters), vol. 22, 143-165, Pergamon Press. Oxford, UK.

Todorov N and Mitev J, 1995. Effect of level of feeding during dry period, and body condition score on reproductive performance in dairy cows, IXth International Conference on Production Diseases in Farm Animals, Sept.11 – 14, Berlin, Germany, p. 302 (Abstr.).


Thesis:

Penkov D, 2008. Estimation of metabolic energy and true digestibility of amino acids of some feeds in experiments with muscovy duck (*Carina moschata*, L). Thesis for DSc. Agrarian University, Plovdiv, 314 pp.

The Editorial Board of the Journal is not responsible for incorrect quotes of reference sources and the relevant violations of copyrights.

AGRICULTURAL SCIENCE AND TECHNOLOGY

Volume 3, Number 4
December 2011



Journal web site:
www.uni-sz.bg/ascitech/index.html


Publisher:
www.alfamarket.biz