



Online Version ISSN: 1314-412X  
Volume 3, Number 4  
December 2011

# *AGRICULTURAL SCIENCE AND TECHNOLOGY*

2011

An International Journal Published by Faculty of Agriculture,  
Trakia University, Stara Zagora, Bulgaria

### **Editor-in-Chief**

*Tsanko Yablanski*  
Faculty of Agriculture  
Trakia University, Stara Zagora  
Bulgaria

### **Co-Editor-in-Chief**

*Radoslav Slavov*  
Faculty of Agriculture  
Trakia University, Stara Zagora  
Bulgaria

### **Editors and Sections**

#### **Genetics and Breeding**

*Atanas Atanassov (Bulgaria)*  
*Ihsan Soysal (Turkey)*  
*Max Rothschild (USA)*  
*Stoitcho Metodiev (Bulgaria)*

#### **Nutrition and Physiology**

*Nikolai Todorov (Bulgaria)*  
*Peter Surai (UK)*  
*Zervas Georgios (Greece)*

#### **Production Systems**

*Dimitar Pavlov (Bulgaria)*  
*Dimitar Panaiotov (Bulgaria)*  
*Jordan Staikov (Bulgaria)*  
*Georgi Zhelyazkov (Bulgaria)*

#### **Agriculture and Environment**

*Georgi Petkov (Bulgaria)*  
*Ramesh Kanwar (USA)*

#### **Product Quality and Safety**

*Marin Kabakchiev (Bulgaria)*  
*Stefan Denev (Bulgaria)*

#### **English Editor**

*Yanka Ivanova (Bulgaria)*

### **Scope and policy of the journal**

Agricultural Science and Technology /AST/ – an International Scientific Journal of Agricultural and Technology Sciences is published in English in one volume of 4 issues per year, as a printed journal and in electronic form. The policy of the journal is to publish original papers, reviews and short communications covering the aspects of agriculture related with life sciences and modern technologies. It will offer opportunities to address the global needs relating to food and environment, health, exploit the technology to provide innovative products and sustainable development. Papers will be considered in aspects of both fundamental and applied science in the areas of Genetics and Breeding, Nutrition and Physiology, Production Systems, Agriculture and Environment and Product Quality and Safety. Other categories closely related to the above topics could be considered by the editors. The detailed information of the journal is available at the website. Proceedings of scientific meetings and conference reports will be considered for special issues.

### **Submission of Manuscripts**

All manuscript written in English should be submitted as MS-Word file attachments via e-mail to [ascitech@uni-sz.bg](mailto:ascitech@uni-sz.bg). Manuscripts must be prepared strictly in accordance with the detailed instructions for authors at the website <http://www.uni-sz.bg/ascitech/index.html> and the instructions on the last page of the journal. For each manuscript the signatures of all authors are needed confirming their consent to publish it and to nominate an author for correspondence. They have to be presented by a submission letter signed by all authors. The form of the submission letter is available upon request from the Technical Assistance or could be downloaded from the website of the journal. All manuscripts are subject to editorial review and the editors reserve the right to improve style and return the paper for rewriting to the authors, if necessary. The editorial board reserves rights to reject manuscripts based on priorities and space availability in the journal.

### **Subscriptions**

Agricultural Science and Technology is published four times a year. The subscription price for institutions is 80 € and for personal subscription 30 € which

include electronic access and delivery. Subscription run for full calendar year. Orders, which must be accompanied by payment may be sent direct to the publisher:

Trakia University  
Faculty of Agriculture, Bank account:  
UniCredit Bulbank,  
Sofia BIC: UNCRBGSF

IBAN: BG29UNCR76303100117681  
With UniCredit Bulbank Stara Zagora

### **Internet Access**

This journal is included in the Trakia University Journals online Service which can be found at [www.uni-sz.bg](http://www.uni-sz.bg).

### **Copyright**


All rights reserved. No part of this publications may be translated into other languages, reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying or any information storage and retrieval system without permission in writing from the publisher.

### **Address of Editorial office:**

Agricultural Science and Technology  
Faculty of Agriculture, Trakia University  
Student's campus, 6000 Stara Zagora  
Bulgaria  
Telephone.: +359 42 699330  
+359 42 699446  
<http://www.uni-sz.bg/ascitech/index.html>

### **Technical Assistance:**

Nely Tzvetanova  
Telephone.: +359 42 699446  
E-mail: [ascitech@uni-sz.bg](mailto:ascitech@uni-sz.bg)



*AGRICULTURAL  
SCIENCE AND TECHNOLOGY*

2011

An International Journal Published by Faculty of Agriculture,  
Trakia University, Stara Zagora, Bulgaria



## Product Quality and Safety

# Fatty acid composition of yogurt supplemented with walnut extract

S. Boycheva<sup>1</sup>, N. Naydenova<sup>2\*</sup>, G. Mihaylova<sup>2</sup>, T. Dimitrov<sup>2</sup>, D. Pavlov<sup>3</sup>

<sup>1</sup>Department of Microbiology, Faculty of Agriculture, Trakia University, 6000 Stara Zagora, Bulgaria

<sup>2</sup>Department of Dairy Science, Faculty of Agriculture, Trakia University, 6000 Stara Zagora, Bulgaria;

<sup>3</sup>Department of Plant Science, Faculty of Agriculture, Trakia University, 6000 Stara Zagora, Bulgaria.

**Abstract.** The present study aimed to monitor the changes in fatty acid composition of yogurt supplemented with ground walnut extract and stored over 10 days. Fatty acid content of raw and pasteurized milk, of 24-hour yogurt and 10-day yogurt (stored at 4°C) was assayed. Total saturated fatty acid content of experimental samples of milk was lower while that of unsaturated fatty acids was higher as compared to control milk, 24-hour and 10-day yogurt samples. Assayed fatty acids exhibited higher differences between control and experimental samples with regard to polyunsaturated (C18:2 and C18:3) fatty acids. The total saturated fatty acid content in pasteurized milk with walnut extract decreased by 11.2% compared to raw milk, while remaining unaltered in natural pasteurized milk. Pasteurized milk with walnuts exhibited higher polyunsaturated fatty acid content – 2.6 times vs natural and almost 3 times higher than raw milk. Yogurt containing walnut extract contained by 19.25% more unsaturated fatty acids and by 2.3 times more polyunsaturated fatty acids as compared to natural yogurt. During storage of produced yogurt for 10 days, the amount of C18:3 decreased by 46.8% in the yogurt containing walnut extract and by 34.4% in natural yogurt.

**Keywords:** fatty acids, yogurt, walnuts

**Keywords:** MUFA – monounsaturated fatty acids, PUFA – polyunsaturated fatty acids, SFA – saturated fatty acids, UFA – unsaturated fatty acids

## Introduction

Milk and dairy products are an essential part of the so-called functional foods. They are aimed at satisfying all dietary needs during the entire human life. Therefore, milk is always a part of human diet and is taking part in the balance of most valuable nutrients. The dietetic and nutritive properties of yogurt make it important part of Bulgarian diet. The considerable rates of consumption of this product are not only a centennial tradition, but are mainly due to its exceptional complete dietary qualities. The functional properties of yogurt could be increased via modification and/or increase of the beneficial effects of its components on human organism. One alternative for modification of Bulgarian yogurt quality (organoleptic and physicochemical parameters) is the supplementation of fruits, seeds, nuts etc. with the purpose to increase its therapeutic, prophylactic or dietetic properties. That is why our attention has been focused on the production of Bulgarian yogurt with supplements beneficial for human health (Dimitrov et al., 2002; Boycheva et al., 2006, 2007; Boycheva et al., 2010, 2011).

During the last years, more and more attention is paid on the fat content of milk and dairy products, including the content of saturated and unsaturated fatty acids. There are two primary classes of polyunsaturated fatty acids –  $\omega$ -6 and  $\omega$ -3, determined by essential fatty acids linoleic and linolenic acids. Both linoleic and linolenic acids are essential, but they could not be synthesized in the human organism. Thus, they should be supplied with food. In previous studies of ours, we monitored the changes in the fatty acids composition of yogurt supplemented with plant extracts (Boycheva et al., 2007a), with walnuts and hazelnuts (Naydenova et al., 2008;

Boycheva et al., 2011).

Health benefits of conjugated linoleic acids as anticarcinogenic (Ha et al., 1990; Ip et al., 1994; Banni et al., 1999; Parodi, 1997; Belury, 2002; Masso-Welch et al., 2002) and antiatherogenic agent (Lee et al., 1994; Pariza et al., 1996; Kritchevsky et al., 2000), immune modulator (Cook et al., 1993; MacDonald, 2000), antidiabetic agent (Schrezenmeir and Jagla, 2000) and a factor reducing the body fat depots (Pariza et al., 1996; Garnsworthy, 1997; DeLany and West, 2000; Blankson et al., 2000) have been demonstrated both *in vitro* and in animal experiments (Pariza, 1999; Hayek et al., 1999; Williams, 2000).

The aim of this research was to monitor the changes in fatty acid composition of yogurt supplemented with ground walnut extract compared to natural yogurt

## Material and methods

### *Fresh milk and starter cultures*

Fresh raw cow milk, obtained from the Experimental Farm of the Trakia University – Stara Zagora and starter culture ready for direct vat inoculation were used for yogurt preparation. The cow's yogurt was prepared in laboratory conditions.

### *Yogurt preparation*

The milk was pasteurized (95°C/30 min), cooled to 45 °C and inoculated with 1.5% yogurt culture consisting of *Lactobacillus delbrueckii ssp. bulgaricus* and *Streptococcus thermophilus*. The raw milk was divided into two lots – control and experimental. The

\* e-mail: n\_naydenova@uni-sz.bg

samples were then cultivated at 42°C until coagulation, then cooled and stored in a refrigerator at 4-6°C. Walnuts (10%) were finely ground and added to milk before pasteurization. After the pasteurization, milk was filtered and they inoculated as described above.

#### Fatty acid composition

The extraction of milk fat was done using the method of Rose-Gottlieb using a diethyl ether and petroleum ether (Methodenbuch, Bd. VI VDLUFA-Verlag, Darmstadt, 1985). After that the solvents were evaporated on a vacuum-rotary evaporator. For obtaining methyl esters of the fatty acids sodium methylate (CHONa) was used (Jahreis et al., 1997). The fatty acid composition of yogurt, both control and experimental, was determined by gas chromatography "Pay-Unicam 304" with flame ionization detector and column EC™-WAX, 30 m, ID 0.25 mm, Film:0,25 µm. The content of the different fatty acid groups and their ratios were calculated. The atherogenic index was calculated according to a formula on the basis of medium-chain fatty acids content – C12:0, C14:0 and C16:0, monounsaturated and polyunsaturated fatty acids (Chilliard et al., 2003).

#### Statistical analysis

For statistical analysis, ANOVA was performed on the two batches and the corresponding replicates, using a statistical software (Statistica 6.0).

## Results and discussion

Data about the fatty acid content of raw and pasteurized milk – natural and supplemented with walnut extract (Table 1) showed that

saturated caproic, caprylic and capric acids decreased in both walnut extract supplemented (by 18.73%, 23.08% and 38.9%, respectively) and in natural pasteurized milk (by 17.31%, 17.31% and 23.24% respectively) compared to raw milk. In a previous study of ours (Boycheva et al., 2007) the concentrations of C8:0 and C10:0 were also lower in pasteurized vs natural milk, but C6:0 content was increased. The experimental batch of pasteurized milk exhibited also reduction in saturated medium-chain fatty acids C12:0, C14:0 and C16:0 as compared to control pasteurized milk. In previous experiments with linseed (Boycheva et al., 2007), lower C12:0 and C14:0 contents in pasteurized milk were also reported.

The content of linoleic acid, an n-6 fatty acid, was almost 3 times higher in pasteurized milk with walnut extract compared to raw milk and 2.4 times higher than that of natural milk. The polyunsaturated linolenic acid was 3 times and 2.7 times higher, respectively, in pasteurized milk with walnut extract compared to natural pasteurized and raw milk. These data are not consistent with previous reports (Mihaylova, 2007) regarding the changes in C18:2, where a reduction of this acid's content has been established.


It is acknowledged that ω-6 and ω-3 fatty acids are precursors of hormone-like substances – eicosanoids, with a primary regulatory role. These molecules possess signalling functions improving the relationships between cells in the nervous and immune systems. They regulate cell growth, inflammatory events, the development of thrombosis, allergic reactions, hypertension etc. (Sarau et al., 1999; Evans et al., 2000; Boyce, 2005). Their effect in most cases is contradictory, that puts an emphasis on the substantial importance of balanced and healthy nutrition.

The total amount of short-chain fatty acids after pasteurization of milk decreased by 23.42% in milk with walnut extract and by 13.69% in natural milk. The total content of saturated fatty acids (SFA) in control milk samples was almost unchanged after the pasteurization, while in pasteurized milk with walnut extract it was by

**Table 1.** Fatty acid content of raw and pasteurized milk

Fatty acids	Raw milk		Pasteurized milk			
			Natural		With walnut extract	
	Mean	SEM	Mean	SEM	Mean	SEM
C 4:0	2.56	0.920	2.73	0.595	2.53	0.112
C 6:0	2.83 <sup>a</sup>	0.441	2.34 <sup>a</sup>	0.243	2.30 <sup>a</sup>	0.375
C 8:0	1.56	0.196	1.29	0.226	1.20	0.211
C 10:0	3.83 <sup>ab</sup>	0.269	2.94 <sup>a</sup>	0.551	2.34 <sup>b</sup>	0.413
C10:1	0.33 <sup>c</sup>	0.036	0.28 <sup>c</sup>	0.065	0.14 <sup>c</sup>	0.045
C 12:0	3.56 <sup>a</sup>	0.389	3.11	0.674	2.58 <sup>a</sup>	0.433
C 14:0	10.26	0.284	10.51	2.362	9.30	1.157
C 14:1	1.07	0.370	0.71 <sup>c</sup>	0.014	1.18 <sup>c</sup>	0.053
C 15 ai	0.40 <sup>a</sup>	0.025	0.51 <sup>a</sup>	0.099	0.45	0.188
C 15:0	0.60	0.184	1.15	0.684	0.48	0.078
C 16:0	33.01 <sup>b</sup>	0.166	33.82 <sup>c</sup>	1.109	27.17 <sup>bc</sup>	2.880
C 18:0	8.52 <sup>b</sup>	1.680	8.85 <sup>b</sup>	0.723	11.26 <sup>b</sup>	0.595
C 18:1	27.06	0.724	26.93	1.712	26.62	1.048
C 18:2	2.75	0.331	3.32	0.742	7.88	2.435
C 18:3	1.67	0.642	1.53	0.325	4.58	1.560

Legend: a – p<0.05, b – p<0.01, c- p<0.001



**Figure 1.** Dynamics of the saturated and unsaturated fatty acids in raw and pasteurized milk

11.2% lower (Figure 1). Total unsaturated fatty acid content (UFA) in pasteurized milk supplemented with walnut extract was by 22.84% higher than that of raw milk and fairly similar to that of natural pasteurized milk. The sums of SFA and UFA in pasteurized milk reported in a previous study were similar (Boycheva et al., 2007).

The sum of monounsaturated fatty acids (MUFA) decreased slightly after the pasteurization of milk in both experimental and control samples, whereas the total content of polyunsaturated fatty acids (PUFA) increased in natural milk samples and was almost 3 times higher in pasteurized milk with walnut extract. The PUFA/SFA


ratio was almost equal in raw and natural pasteurized milk (0.06 and 0.07, respectively), whereas in pasteurized milk supplemented with walnut extract it was higher – 0.21 by reason of the higher UFA concentration. The  $\omega$ -6/  $\omega$ -3 ratio was balanced from the point of view of healthy nutrition – from 1.65 to 2.17, in raw and pasteurized milk samples, respectively. The atherogenic index was the lowest in experimental pasteurized milk – 1.66 vs 2.36–2.41 for raw and pasteurized natural milk.

The fatty acid composition of the product is presented in Table 2. The content of the individual short-chain fatty acids changed slightly

**Table 2.** Fatty acids in yogurt

Fatty acids	24-hour yogurt			
	Natural		With walnut extract	
	Mean	SEM	Mean	SEM
C 4:0	2.69	0.380	2.68	0.012
C 6:0	2.57	0.279	2.30	0.226
C 8:0	1.22	0.031	1.18	0.078
C 10:0	2.79	0.384	2.85	0.042
C 12:0	2.94	0.205	2.66	0.057
C 13:0	0.18	0.070	-	-
C 14:0	10.75	0.951	9.91	0.559
C 14:1	0.69 <sup>e</sup>	0.094	1.16 <sup>e</sup>	0.018
C 15:0	1.02 <sup>e</sup>	0.045	0.38 <sup>e</sup>	0.003
C 16:0	34.94 <sup>b</sup>	1.998	29.08 <sup>b</sup>	3.401
C 18:0	8.23	0.857	10.00	0.276
C 18:1	26.62	3.435	25.60	0.969
C 18:2	3.55	1.697	7.56	0.231
C 18:3	1.83 <sup>b</sup>	0.585	4.66 <sup>b</sup>	0.488

Legend: a –  $p < 0.05$ , b –  $p < 0.01$ , c –  $p < 0.001$



**Figure 2.** Dynamics of saturated and unsaturated fatty acids in control and experimental yogurt at 24 h

after the lactic acid fermentation while the concentrations of some medium- and long-chain fatty acids in yogurt supplemented with walnut extract exhibited various trends is the natural yogurt. The caproic acid in walnut yogurt was by 10.5% lower compared to control yogurt. The changes in caprylic and capric acids were insignificant. The contents of long-chain fatty acids – C18:2 and C18:3 were higher in milk with walnut extract compared to natural yogurt, unlike the concentration of monounsaturated C18:1. Lower oleic acid content in experimental samples was probably due to lactic acid bacteria and more specifically, to lactobacilli which need this acid and whose counts in this yogurt sample were higher, as also

shown in previous studies (Naydenova et al., 2008). The amounts of essential linoleic and linolenic acid in yogurt supplemented with walnut extract were significantly higher (2.13 and 2.55 times respectively) than in control samples.

Yogurt with walnut extract exhibited the same tendency as pasteurized milk – increased total PUFA amount (2.3 times) and slightly lower MUFA content. The UFA amount was higher (by 19.25%) whereas SFA decreased by 9.34% in yogurt produced from walnut extract-supplemented milk (Figure 2). The PUFA/SFA ratio in walnut yogurt was 2.5 times higher than that of natural yogurt. The atherogenic index and the n-6/n-3 ratio were lower.

**Table 3.** Fatty acids in yogurt after storage for 10 days


Fatty acids	10-day yogurt			
	Natural		With walnut extract	
	Mean	SEM	Mean	SEM
C 4:0	2.29	0.135	2.40	0.226
C 6:0	2.48	0.138	2.47	0.035
C 8:0	1.39	0.106	1.37	0.071
C 10:0	3.23	0.333	2.99	0.318
C10:1	0.32 <sup>c</sup>	0.008	3.08 <sup>c</sup>	0.332
C 12:0	3.30	0.255	2.89	0.339
C 12:1	0.08 <sup>c</sup>	0.001	0.66 <sup>c</sup>	0.063
C 14:0	10.74	0.485	9.78	0.297
C 14:1	0.76	0.020	1.01	0.085
C 15:0	0.91	0.123	0.94	0.057
C 16:0	35.23 <sup>c</sup>	0.361	27.41 <sup>c</sup>	0.940
C 18:0	8.24	1.134	10.03	0.827
C 18:1	26.35	5.408	25.20	2.135
C 18:2	3.48 <sup>c</sup>	0.418	7.32 <sup>c</sup>	0.204
C 18:3	1.2 <sup>b</sup>	0.002	2.48 <sup>b</sup>	0.025

Legend: a – p<0.05; b – p<0.01; c- p<0.001



During storage in a refrigerator for 10 days (Table 3), no significant changes in individual fatty acids have occurred in either natural or walnut extract supplemented yogurt with the exception of C16:0 and C18:3 contents in experimental samples. The amount of C18:3 decreased by 46.8% during storage in experimental samples and by 34.4% in control samples as compared to respective levels in 24-hour yogurt. The ratio C18:2/C18:3 increased in both yogurts to

almost equal levels – 2.9 in natural yogurt and 2.95 in walnut yogurt. The PUFA/SFA ratios decreased after 10 days of storage compared to 24-hour storage to 0.07 in control and 1.16 in experimental samples. The total PUFA content decreased during the storage. The 10-day storage of yogurt containing walnut extract resulted in slight reduction of saturated fatty acids and increased unsaturated fatty acids, mainly on account of monounsaturated fatty acids (Figure 3.).



**Figure 3.** Dynamics of the saturated and unsaturated fatty acids in control and experimental yogurt at 10 days

## Conclusion

The total saturated fatty acid content in pasteurized milk with walnut extract decreased by 11.2% compared to raw milk, while remaining unaltered in natural pasteurized milk. Pasteurized milk with walnuts exhibited higher polyunsaturated fatty acid content – 2.6 times vs natural and almost 3 times higher than raw milk. Yogurt containing walnut extract contained by 19.25% more unsaturated fatty acids and by 2.3 times more polyunsaturated fatty acids as compared to natural yogurt. During storage of produced yogurt for 10 days, the amount of C18:3 decreased by 46.8% in the yogurt containing walnut extract and by 34.4% in natural yogurt.

## References

**Boycheva S, Dimitrov T, Mihaylova G and Pavlov D**, 2006. Effect of some plant supplements on fatty acid content of Bulgarian yogurt. Proceedings from the 1st International Symposium "Ecological approaches in safe foods production", Plovdiv, October 19-20, 351-356.

**Dimitrov T, Boycheva S, Iliev T and Zheleva N**, 2002. Production of fruit yogurt with different supplements. Proceedings of the Scientific Conference with International Participation, Union of Scientists, Bulgaria, 2, 323-327(Bg).

**Mihaylova G**, 2007. Fatty acid composition and functional properties of the fat in milk and milk products from different agricultural ecosystems, Thesis for DSc, Trakia University, Stara

Zagora.

**Naydenova N, Boycheva S, Mihaylova G, Dimitrov T and Pavlov D**, 2008. Yogurt supplemented with walnut extract. *Animal Science*, 6, 57-62.

**Banni S, Angioni A, Casu V, Melis MP, Carta G, Corongiu F, Thompson H and Ip C**, 1999. Decrease in linoleic acid metabolites as a potential mechanism in cancer risk reduction by conjugated linoleic acid. *Carcinogenesis*, 20, 6, 1019-1024.

**Belury MA**, 2002. Inhibition of carcinogenesis by conjugated linoleic acid: Potential mechanisms of action. *Journal of Nutrition*, 132, 10, 2995-2998.

**Blankson H, Stakkestad JA, Fagertun H, Thom E, Wadstein J and Gudmundsen O**, 2000. Conjugated linoleic acid reduces body fat mass in overweight and obese humans. *Journal of Nutrition*, 130, 12, 2943-2948.

**Boycheva S, Dimitrov T, Mihaylova G and Pavlov D**, 2007. Effect of some plant supplements on the quality of fatty acid and mineral elements in Bulgarian fermented milk. *Food processing industry magazine – Specializing edition Food Science*, 2, 2-7.

**Boycheva S, Dimitrov T, Mihaylova G and Naydenova N**, 2007a. Aminoacids, fatty acids and mineral composition of yogurt made from cow's and goat's milk added with probiotic bacteria. *Food Science*, 5, 39-41.

**Boycheva S, Naydenova N, Mihaylova G, Dimitrov T and Pavlov D**, 2010. Quality characteristics of yogurt supplemented with nuts. *Agricultural Science and Technology*, 4, 221-226.

**Boycheva S, Dimitrov T, Naydenova N and Mihaylova G**, 2011. Quality characteristics of yogurt from goat's milk, supplemented with

fruit juice. *Czech Journal of Food Science*, 1, 24-30.

**Boyce J**, 2005. Eicosanoid mediators of mast cells: receptors, regulation of synthesis and pathobiologic implications. *Chemical Immunology and Allergy*, 87, 59-79.

**Chilliard Y, Ferlay A, Rouel J and Lamberet G**, 2003. A review of nutritional and physiological factors affecting goat milk synthesis and lipolysis. *Journal of Dairy Science*, 86, 1751-1770.

**Cook ME, Miller CC, Park Y and Pariza MW**, 1993. Immune modulation by altered nutrient metabolism: nutritional control of immune-induced growth depression. *Poultry Science*, 72, 1301-1305.

**DeLany JP and West DB**, 2000. Changes in body composition with conjugated linoleic acid. *Journal of the American College of Nutrition*, 19, 4, 487S-493.

**Dimitrov T, Boycheva S, Iliev T and Zheleva N**, 2002. Production of yogurt with different additives. *Animal Science*, 4-5, 41-44.

**Evans AR, Junger H and Southall MD**, 2000. Isoprostanes, novel eicosanoids that produce nociception and sensitize rat sensory neurons. *The Journal of Pharmacology and Experimental Therapeutics*, 293, 3, 912-920.

**Jahreis G, Fritsche J and Steinhart H**, 1997. Conjugated linoleic acid in milk fat: high variation depending on production system. *Nutrition Research*, 17, 9, 1479-1484.

**Garnsworthy PC**, 1997. Fat in dairy cows. In: *Recent advances in animal nutrition*, Garnsworthy PC and Wiseman J, ed. Nottingham University Press. Nottingham, UK, 87-104.

**Ha YL, Storkson J and Pariza MW**, 1990. Inhibition of benzo (a) pyrene-induced mouse forestomach neoplasia by conjugated dienoic derivatives of linoleic acid. *Cancer research*, 50, 1097-1101.

**Hayek MG, Han SN, Wu D, Watkins BA, Meydani M, Dorsey JL, Smith DE and Meydani SN**, 1999. Dietary conjugated linoleic acid influences the immune response of young and old C57BL/6NCrIBR mice. *Journal of Nutrition*, 129, 1, 32-38.

**Ip C, Singh M, Thompson HJ and Scimeca J**, 1994. Conjugated

linoleic acid suppresses mammary carcinogenesis and proliferative activity of the mammary gland in the rat. *Cancer Research*, 54, 5, 1212-1215.

**Kritchevsky D, Tepper SA, Wright S, Tso P and Czarnecki SK**, 2000. Influence of conjugated linoleic acid (CLA) on establishment and progression of atherosclerosis in rabbits. *Journal of the American College of Nutrition*, 19, 4, 472S-477.

**Lee KN, Kritchevsky D and Pariza MW**, 1994. Conjugated linoleic acid and atherosclerosis in rabbits. *Atherosclerosis*, 108, 19-25.

**MacDonald HB**, 2000. Conjugated linoleic acid and disease prevention: A review of current knowledge. *Journal of the American College of Nutrition*, 19, 111S-118.

**Masso-Welch PA, Zangani D, Ip C, Vaughan MM, Shoemaker S, Ramirez RA and Ip MM**, 2002. Inhibition of angiogenesis by the cancer chemopreventive agent conjugated linoleic acid. *Cancer Research*, 62, 15, 4383-4389.

**Pariza MW**, 1999. The biological activities of conjugated linoleic acid. Ch. 2 In: Yuraweca MP, Mossoba MM, Kramer JKG, Pariza MW, Nelson GJ ed. *Advances in conjugated linoleic acid research*, 1, Champaign, Ill. AOCs Press.

**Pariza MW, Park Y, Cook M, Albright K and Liu W**, 1996. Conjugated linoleic acid (CLA) reduces body fat. *FASEB Journal*, 10, 3227.

**Parodi PW**, 1997. Cow's milk fat components as potential anticarcinogenic agents. *Journal of Nutrition*, 127, 6, 1055-1060.

**Sarau HM, Foley JJ and Schmidt DB**, 1999. In vitro and in vivo pharmacological characterization of SB 201993, an eicosanoid-like LTB<sub>4</sub> receptor antagonist with anti-inflammatory activity. *Prostaglandins, Leukotrienes and Essential Fatty Acids*, 61, 1, 55-64.

**Schrezenmeir J and Jagla A**, 2000. Milk and diabetes. *Journal of the American College of Nutrition*, 19, 176S-190.

**William C**, 2000. Commentary: is CLA an important dietary variable relevant to chronic disease of humans. *Nutrition Bulletin*, 25, 29.

**CONTENTS**

1 / 2

**Reviews**

---

- 1. Lameness scoring systems for cattle in dairy farms** 291  
T. Penev

**Genetics and Breeding**

---

- 2. Body and carcass parameters of sea bream (*Sparus aurata* L.) and sea bass (*Dicentrarchus labrax* L.)** 299  
M. Marinova, I. Sirakov, Y. Staykov, E. Ivancheva
- 3. Effect of breed upon blood lysozyme and complement activity in different sheep breeds** 302  
L. Sotirov, Ts. Koynarski, V. Semerdjiev, D. Dimov, S. Laleva, P. Slavova, M. Iliev, D. Yarkov
- 4. Winter wheat productivity under favorable and drought environments III. Effect of fertilization** 306  
A. Ivanova, N. Tsenov
- 5. Evaluation of perspective sorghum breeding forms in their reaction to some diseases in field conditions** 310  
M. Georgieva – Andreeva, K. Tanova, S. Raykov

**Nutrition and Physiology**

---

- Effect of dietary coconut oil supplementation on some blood biochemical indices in yearling rams** 313  
T. Slavov, V. Radev, K. Sivkova, I. Varlyakov
- Pharmacokinetics of tilmicosin after oral application of Pulmotil G 200 – premix in pigs** 318  
D. Dimitrova, V. Katsarov, D. Dimitrov, D. Tsoneva
- Epidermal growth factor content in rabbit doe milk during the different lactation stages** 323  
E. Vachkova, B. Bivolarski
- Intraorbital glands in turkey broilers. III. Lacrimal gland histometry** 327  
D. Dimitrov
- Effect of body condition score at calving on body condition during lactation in Holstein and Brown Swiss cows** 330  
Zh. Gergovska, T. Angelova, D. Yordanova, Zh. Krastanov, Ch. Miteva
- Use of brewer's grains for feeding of lambs** 336  
A. Kirilov, K. Ivanov

**Production Systems**

---

- The effect of the milking liner design on the parameters of the milking machine pulsation system** 339  
V. Vlashev, B. Banev, K. Peichev, G. Dineva
- Accumulation dynamic of *Ruta graveolens* L. essential oil** 343  
A. Dzurmanski, G. Zhekova, D. Angelova
- Research on the water regimen of soil upon the production of vine planting material** 346  
N. Kovachev, N. Taneva, V. Kovachev, L. Halil

**Agriculture and Environment**

---

- Influence of the farming, soil cultivation and Fertilization on the yield of wheat** 351  
M. Nankov, L. Glogova
- Study on the applicability of a natural geomaterial for mononitrophenol removal from simulated agricultural run-off water** 354  
Z. Yaneva, B. Koumanova, N. Georgieva

**CONTENTS**

2 / 2

---

<b>Comparative technical and economic analysis of variants for cleaning and storage of manure on a farm for 108 – 120 dairy cows</b>	<b>359</b>
V. Dimova, D. Dinev, Y. Popova, Y. Mitev	
<b>Distribution of the black mussel <i>Mytilus galloprovincialis</i> (L.) along the Bulgarian Black Sea coast</b>	<b>368</b>
E. Petrova, St. Stoykov	
<b>Toxicity of plant protection products towards the imago of <i>Encarsia Formosa</i> Gah.</b>	<b>374</b>
V. Yankova, S. Masheva, B. Boev, K. Toskov	
<b>Effect of the rhizobacterium <i>Bacillus subtilis</i> on the development of the root-knot nematode <i>Meloidogyne arenaria</i> at different temperatures</b>	<b>378</b>
M. Mohamedova, H. Samaliev	
<b>Product Quality and Safety</b>	
<hr/>	
<b>Fatty acid composition of yogurt supplemented with walnut extract</b>	<b>384</b>
S. Boycheva, N. Naydenova, G. Mihaylova, T. Dimitrov, D. Pavlov	
<b>Near Infrared Spectroscopy for monitoring changes during yellow cheese ripening</b>	<b>390</b>
S. Atanassova, N. Naydenova, T. Kolev, T. Iliev, G. Mihaylova	
<b>Short communications</b>	
<hr/>	
<b>Mechanical correction the traction weight of a farm wheeled tractor</b>	<b>395</b>
D. Irinchev	
<b>Possibilities for increasing the yield and quality of asters (<i>Callistephus chinensis</i> L.) cut flower</b>	<b>397</b>
N. Miteva, O. Tafradziiski	

## **Instruction for authors**

### **Preparation of papers**

Papers shall be submitted at the editorial office typed on standard typing pages (A4, 30 lines per page, 62 characters per line). The editors recommend up to 15 pages for full research paper (including abstract, references, tables, figures and other appendices)

**The manuscript** should be structured as follows: Title, Names of authors and affiliation address, Abstract, List of keywords, Introduction, Material and methods, Results, Discussion, Conclusion, Acknowledgements (if any), References, Tables, Figures.

**The title** needs to be as concise and informative about the nature of research. It should be written with small letter /bold, 14/ without any abbreviations.

### **Names and affiliation of authors**

The names of the authors should be presented from the initials of first names followed by the family names. The complete address and name of the institution should be stated next. The affiliation of authors are designated by different signs. For the author who is going to be corresponding by the editorial board and readers, an E-mail address and telephone number should be presented as footnote on the first page. Corresponding author is indicated with \*.

**Abstract** should be not more than 350 words. It should be clearly stated what new findings have been made in the course of research. Abbreviations and references to authors are inadmissible in the summary. It should be understandable without having read the paper and should be in one paragraph.

**Keywords:** Up to maximum of 5 keywords should be selected not repeating the title but giving the essence of study.

**The introduction** must answer the following questions: What is known and what is new on the studied issue? What necessitated the research problem, described in the paper? What is your hypothesis and goal?

**Material and methods:** The objects of research, organization of experiments, chemical analyses, statistical and other methods and conditions applied for the experiments should be described in detail. A criterion of sufficient information is to be

possible for others to repeat the experiment in order to verify results.

**Results** are presented in understandable tables and figures, accompanied by the statistical parameters needed for the evaluation. Data from tables and figures should not be repeated in the text.

**Tables** should be as simple and as few as possible. Each table should have its own explanatory title and to be typed on a separate page. They should be outside the main body of the text and an indication should be given where it should be inserted.

**Figures** should be sharp with good contrast and rendition. Graphic materials should be preferred. Photographs to be appropriate for printing. Illustrations are supplied in colour as an exception after special agreement with the editorial board and possible payment of extra costs. The figures are to be each in a single file and their location should be given within the text.

**Discussion:** The objective of this section is to indicate the scientific significance of the study. By comparing the results and conclusions of other scientists the contribution of the study for expanding or modifying existing knowledge is pointed out clearly and convincingly to the reader.

**Conclusion:** The most important consequences for the science and practice resulting from the conducted research should be summarized in a few sentences. The conclusions shouldn't be numbered and no new paragraphs be used. Contributions are the core of conclusions.

### **References:**

In the text, references should be cited as follows: single author: Sandberg (2002); two authors: Andersson and Georges (2004); more than two authors: Andersson et al. (2003). When several references are cited simultaneously, they should be ranked by chronological order e.g.: (Sandberg, 2002; Andersson et al., 2003; Andersson and Georges, 2004). References are arranged alphabetically by the name of the first author. If an author is cited more than once, first his individual publications are given ranked by year, then come publications with one co-author, two co-authors, etc. The names of authors, article and journal titles in the Cyrillic or alphabet different from Latin, should be transliterated into Latin and article titles should be translated into English. The original language of articles and books translated into English is indicated in

parenthesis after the bibliographic reference (Bulgarian = Bg, Russian = Ru, Serbian = Sr, if in the Cyrillic, Mongolian = Mo, Greek = Gr, Georgian = Geor., Japanese = Ja, Chinese = Ch, Arabic = Ar, etc.)

The following order in the reference list is recommended:

**Journal articles:** Author(s) surname and initials, year. Title. Full title of the journal, volume, pages. Example:

**Simm G, Lewis RM, Grundy B and Dingwall WS**, 2002. Responses to selection for lean growth in sheep. *Animal Science*, 74, 39-50

**Books:** Author(s) surname and initials, year. Title. Edition, name of publisher, place of publication. Example: **Oldenbroek JK**, 1999. Genebanks and the conservation of farm animal genetic resources, Second edition. DLO Institute for Animal Science and Health, Netherlands.

**Book chapter or conference proceedings:** Author(s) surname and initials, year. Title. In: Title of the book or of the proceedings followed by the editor(s), volume, pages. Name of publisher, place of publication. Example:

**Mauff G, Pulverer G, Operkuch W, Hummel K and Hidden C**, 1995. C3-variants and diverse phenotypes of unconverted and converted C3. In: *Provides of the Biological Fluids* (ed. H. Peters), vol. 22, 143-165, Pergamon Press. Oxford, UK.

**Todorov N and Mitev J**, 1995. Effect of level of feeding during dry period, and body condition score on reproductive performance in dairy cows, IX<sup>th</sup> International Conference on Production Diseases in Farm Animals, Sept. 11 – 14, Berlin, Germany, p. 302 (Abstr.).

### **Thesis:**

**Penkov D**, 2008. Estimation of metabolic energy and true digestibility of amino acids of some feeds in experiments with muscovy duck (*Carina moschata*, L). Thesis for DSc. Agrarian University, Plovdiv, 314 pp.

The Editorial Board of the Journal is not responsible for incorrect quotes of reference sources and the relevant violations of copyrights.

# AGRICULTURAL SCIENCE AND TECHNOLOGY

Volume 3, Number 4  
December 2011



Journal web site:  
[www.uni-sz.bg/ascitech/index.html](http://www.uni-sz.bg/ascitech/index.html)

  
Publisher:  
[www.alfamarket.biz](http://www.alfamarket.biz)